

Recombinant human CD40 Ligand/CD40LG protein

Catalog Number: ATGP4019

PRODUCT INFORMATION

Expression system

E.coli

Domain

113-261aa

UniProt No.

P29965

NCBI Accession No.

NP_000065

Alternative Names

CD40 ligand, CD40-L, CD40L, T-cell antigen Gp39, TNF-related activation protein, TRAP, Tumor necrosis factor ligand superfamily member 5, TNFSF5, CD154, hyper-IgM syndrome, HIGM1, T-B cell-activating molecule, T-BAM

PRODUCT SPECIFICATION

Molecular Weight

18.3 kDa (169aa) confirmed by MALDI-TOF

Concentration

0.5mg/ml (determined by Bradford assay)

Formulation

Liquid in. 20mM Tris-HCl buffer (pH 8.0) containing 1mM DTT, 40% glycerol, 0.2M NaCl

Purity

> 95% by SDS-PAGE

Endotoxin level

< 1 EU per 1ug of protein (determined by LAL method)

Biological Activity

Measured by its binding ability in a functional ELISA with Human CD40/TNFRSF5 (CAT# ATGP4007).

Tag

His-Tag

Application

SDS-PAGE, Bioactivity

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Recombinant human CD40 Ligand/CD40LG protein

Catalog Number: ATGP4019

Description

CD40 ligand, also known as CD40LG, is a member of the TNF superfamily and is expressed on activated T cells. CD40LG is expressed as a soluble cytokine as well as a homotrimeric type II transmembrane protein. It has been reported to be important for B cell costimulation following binding of its receptor, CD40. Mutations in the gene encoding for CD154 are implicated in hyper-IgM immunodeficiency syndrome type 1. Recombinant human CD40LG protein, fused to His-tag at N-terminus, was expressed in E. coli and purified by using conventional chromatography techniques.

Amino acid Sequence


<MGSSHHHHHH SGLVPRGSH> MQKGDQNPQI AAHVISEASS KTTSVLQWAE KGYTMSNNL VTLENGKQLT VKRQGLYYIY AQTFCSNRE ASSQAPFIAS LCLKSPGRFE RILLRAANTH SSAKPCGQQS IHLGGVFELQ PGASVFNVT DPSQVSHGTG FTSFGLLKL

General References

Gordon J., et al. (1995) Blood Rev. 9:53-56.
Cheng G., et al. (1995) Science. 267:1494-1498.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

Biological Activity

Human CD40 Ligand/TNFSF5 is coated at 5 ug/ml (100 ul/well) can bind Human CD40/TNFRSF5 (CAT# ATGP4007) in a Functional ELISA assay.

