NKMAXBio We support you, we believe in your research

Recombinant canine IL-8/CXCL8 protein

Catalog Number: ATGP4043

PRODUCT INFORMATION

Expression system

HEK293

Domain

28-101aa

UniProt No.

P41324

NCBI Accession No.

NP 001003200.1

Alternative Names

Interleukin-8, C-X-C motif chemokine 8, Chemokine (C-X-C motif) ligand 8

PRODUCT SPECIFICATION

Molecular Weight

9.4 kDa (80aa)

Concentration

0.25mg/ml (determined by BCA assay)

Formulation

Liquid. In Phosphate-Buffered Saline (pH 7.4) containing 20% glycerol

Purity

> 90% by SDS-PAGE

Endotoxin level

< 1 EU per 1ug of protein (determined by LAL method)

Tag

His-Tag

Application

SDS-PAGE

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

IL-8/CXCL8, also known as Interleukin-8, GCP-1, and NAP-1 is a member of the CXC family of chemokines family which is widely expressed proinflammatory. IL-8 is a potent neutrophil chemotactic factor. It attracts neutrophils, basophils, and T-cells, but not monocytes. This protein has two primary functions. It induces chemotaxis in target cells, primarily neutrophils but also other granulocytes, causing them to migrate toward the site of



NKMAXBio We support you, we believe in your research

Recombinant canine IL-8/CXCL8 protein

Catalog Number: ATGP4043

infection. IL-8 also stimulates phagocytosis once they have arrived. IL-8 is also known to be a potent promoter of angiogenesis. In target cells, IL-8 induces a series of physiological responses required for migration and phagocytosis, such as increases in intracellular Ca2+, exocytosis, and the respiratory burst. Among the eceptors on the surface membrane capable of binding IL-8, the most frequently studied receptors are CXCR1 and CXCR2. Expression and affinity for IL-8 differs between the two receptors (CXCR1 > CXCR2). Recombinant canine IL-8/CXCL8, fused to His-tag at C-terminus, was expressed in HEK293 cell and purified by using conventional chromatography techniques.

Amino acid Sequence


VSSELRCQCI KTHSTPFHPK YIKELRVIDS GPHCENSEII VKLFNGNEVC LDPKEKWVQK VVQIFLKKAE KQDP<HHHHHHH>

General References

Lazennec, G. and A. Richmond (2010) Trends Mol. Med. 16:133-144. Munoz, L.M. et al. (2009) J. Immunol. 183:7337-7346. Bergin, D.A. et al. (2010) J. Clin. Invest. 120:4236-4250.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain

