NKMAXBio We support you, we believe in your research

Recombinant canine IL-17A protein

Catalog Number: ATGP4068

PRODUCT INFORMATION

Expression system

HEK293

Domain

29-155aa

UniProt No.

C6L8D7

NCBI Accession No.

NP 001159350.1

Alternative Names

IL17A, IL17, IL-17A, Interleukin 17A, Interleukin-17A, cytotoxic T-lymphocyte-associated serine esterase 8, CTLA-8, CTLA8

PRODUCT SPECIFICATION

Molecular Weight

15.6kDa (133aa)

Concentration

0.5 mg/ml (determined by Absorbance at 280nm)

Formulation

Liquid. In Phosphate-Buffered Saline (pH 7.4) containing 20% glycerol.

Purity

> 90% by SDS - PAGE

Endotoxin level

< 1 EU per 1ug of protein (determined by LAL method)

Biological Activity

The activity is determined by the IL-6 ELISA in a using NIH/3T3 mouse embryonic fibroblast cells. The ED50 range \leq 10 ng/ml.

Tag

His-Tag

Application

SDS-PAGE, Bioactivity

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.



NKMAXBIO We support you, we believe in your research

Recombinant canine IL-17A protein

Catalog Number: ATGP4068

BACKGROUND

Description

IL-17A, also known as CTLA8, is a member of the IL-17 family of cytokines. This family demonstrate a structural motif termed a cysteine knot that also characterizes a large superfamily of growth factors. The IL-17 family of cytokines includes six members, IL-17A, IL-17B, IL-17C, IL-17D, IL-17E/IL-25, and IL-17F. IL-17A main funcion is stimulates production of IL-6, prostaglandin E and nitric oxide. And other inflammatory cytokines(TNF-alpha, IL-1 beta and IFN -gamma) leads to up-regulation of gene expression and progression and amplification of local inflammation. In addition, it serves induction of co-stimulatory molecules such as ICAM-1 and other cytokines, thereby supporting T cell activation. IL-17 expression has been associated with several chronic inflammatory diseases including rheumatoid arthritis, psoriasis and multiple sclerosis. Recombinant canine IL-17A, fused to Histag at C-terminus, was expressed in HEK293 cell and purified by using conventional chromatography techniques.

Amino acid Sequence


FPQNPGCRNT EDKNFPQHVK VNLNILNRNT NSRRPSDYYN RSTSPWNLHR NEDPERYPSV IWEAKCRHLG CVNNEGNINY HMNSVPIQQE ILVLRRESQH CPHSFRLEKM LVAVGCTCVT PIVRHVA<HHH HHH>

General References

Moseley TA., et al. (2003) Cytokine & Growth Factor Reviews. 14: 155–174. Afzali, B. et al. (2007) Clin. Exp. Immunol. 148:32-46. Fossiez, F. et al. (1996) J. Exp. Med. 183:2593-2603.

DATA

SDS-PAGE




3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain

Biological Activity

NKMAXBio We support you, we believe in your research

Recombinant canine IL-17A protein

Catalog Number: ATGP4068



Canine IL-17A stimulates IL-6 secretion of the NIH/3T3 mouse embryonic fibroblast cells. The ED50 range \leq 10 ng/ml.

