NKMAXBIO We support you, we believe in your research

Recombinant human TREM2 protein

Catalog Number: ATGP4104

PRODUCT INFORMATION

Expression system

HEK293

Domain

19-174aa

UniProt No.

O9NZC2

NCBI Accession No.

NP 061838.1

Alternative Names

triggering receptor expressed on myeloid cells 2 isoform 1, triggering receptor expressed on myeloid cells 2, triggering receptor expressed on myeloid cells 2a, TREM-2, Trem2a, Trem2b, Trem2c

PRODUCT SPECIFICATION

Molecular Weight

18.2kDa (162aa)

Concentration

1mg/ml (determined by absorbance at 280nm)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity

> 95% by SDS-PAGE

Endotoxin level

< 1 EU per 1ug of protein (determined by LAL method)

Tag

His-Tag

Application

SDS-PAGE

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

TREM2, also known as Triggering receptor expressed on myeloid cells 2 isoform 1, is a membrane protein that forms a receptorsignaling complex with the TYRO protein tyrosine kinase binding protein. This protein is found in immune cells termed myeloid cells, which include macrophages, granulocytes, monocytes, and dendritic cells. It



NKMAXBio We support you, we believe in your research

Recombinant human TREM2 protein

Catalog Number: ATGP4104

functions in immune response and maybe involved in chronic inflammation by triggering the production of constitutive inflammatory cytokines. It also promotes the differentiation and function of osteoclasts, the production of inflammatory cytokines by adipocytes, insulin resistance, and the phagocytic clearance of bacteria. Defects in TREM2 are acause of polycystic lipomembranous osteodysplasia with sclerosingleukoencephalopathy (PLOSL). In addition, TREM2 is overexpressed in many tumor types and has anti-inflammatory activities. It might therefore be a good therapeutic target. Recombinant human TREM2, fused to His-tag at C-terminus, was expressed in HEK293 cell and purified by using conventional chromatography techniques.

Amino acid Sequence


HNTTVFQGVA GQSLQVSCPY DSMKHWGRRK AWCRQLGEKG PCQRVVSTHN LWLLSFLRRW NGSTAITDDT LGGTLTITLR NLQPHDAGLY QCQSLHGSEA DTLRKVLVEV LADPLDHRDA GDLWFPGESE SFEDAHVEHS ISRSLLEGEI PFPPTS<HHHH HH>

General References

Bouchon A., et al (2000). J. Immunol. 164:4991-4995. N' Diaye EN, et al.(2009). J Cell Biol. 184:215-223.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain

