

Recombinant human TrkB protein

Catalog Number: ATGP4152

PRODUCT INFORMATION

Expression system

HEK293

Domain

32-430aa

UniProt No.

Q16620

NCBI Accession No.

NP_001018074.1

Alternative Names

GP145-TrkB, trk-B, TRKB, NTRK2, OBHD, BDNF/NT-3 growth factors receptor isoform c, BDNF/NT-3 growth factors receptor, Neurotrophic tyrosine kinase receptor type 2, TrkB tyrosine kinase, Tropomyosin-related kinase B

PRODUCT SPECIFICATION

Molecular Weight

70.3kDa (632aa)

Concentration

1mg/ml (determined by absorbance at 280nm)

Formulation

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

Purity

> 90% by SDS-PAGE

Endotoxin level

< 1 EU per 1ug of protein (determined by LAL method)

Tag

hIgG-Tag

Application

SDS-PAGE

Storage Condition

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

BACKGROUND

Description

TrkB/NTRK2, also known as BDNF/NT-3 growth factors receptor, is a receptor tyrosine kinase involved in the development and the maturation of the central and the peripheral nervous systems through regulation of neuron survival, proliferation, migration, differentiation, and synapse formation and plasticity. Trk family are four

Recombinant human TrkB protein

Catalog Number: ATGP4152

members ; TrkA, TrkB, TrkC and a related p75NTR receptor. Each family member binds different neurotrophins with varying affinities and TrkB has the highest affinity for BDNF. It plays a role in learning and memory by regulating both short term synaptic function and long-term potentiation. Mutations in TrkB have been associated with obesity and mood disorders. Recombinant human TrkB/NTRK2, fused to hlgG-tag at C-terminus, was expressed in HEK293 cell and purified by using conventional chromatography techniques.

Amino acid Sequence


CPTSCKCSAS RIWCSDPSPG IVAFPRLEPN SVDPENITEI FIANQKRLEI INEDDVEAYV GLRNLTIVDS GLKFVAHKAF
LKNSNLQHIN FTRNKLTSLR RKHFRHLDLS ELILVGNPFT CSCDIMWIKT LQEAQSSPDT QDLYCLNESS KNIPLANLQI
PNCGLPSANL AAPNLTVEEG KSITLSCSVA GDPVPMYWD VGNLVSKHMN ETSHTQGLR ITNISSDDSG KQISCVAENL
VGEDQDSVNL TVHFAPTITF LESPTSDHHW CIPFTVKGNP KPALQWFYNG AILNESKYIC TKIHVTNHTE YHGCLQLDNP
THMNNGDYTL IAKNEYGKDE KQISAHFMGW PGIDDGANPN YPDVIYEDYG TAANDIGDIT NRSNEIPSTD VTDKTGREH<L
EPKSCDKTHT CPPCPAPELL GGPSVFLFPP KPKDTLMISR TPEVTCVVVD VSHEDPEVKF NWYVDGVEVH NAKTKPREEQ
YNSTYRVVSV LTVLHQDWLN GKEYKCKVSN KALPAIEKT ISKAKGQPRE PQVYTLPPSR DELTKNQVSL TCLVKGFYPS
DIAVEWESNG QPENNYKTP PVLDSGDSFF LYSKLTVDKS RWQQGNVFSC SVMHEALHNNH YTKSLSLSP GK>

General References

Yeo GS., et al. (2004) Nat Neurosci. 7:1187-1189.
Banfield MJ., et al. (2001) Structure. 9:1191-1199.

DATA

SDS-PAGE



3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.