# NKMAXBio We support you, we believe in your research

# Recombinant human CD23/FCER2 protein

Catalog Number: ATGP4162

### **PRODUCT INFORMATION**

# **Expression system**

Baculovirus

#### **Domain**

48-321aa

#### UniProt No.

P06734

#### **NCBI Accession No.**

NP 001993.2

#### **Alternative Names**

Low affinity immunoglobulin epsilon Fc receptor, BLAST-2, C-type lectin domain family 4 member J, Fc-epsilon RII, Immunoglobulin E-binding factor, Lymphocyte IgE receptor, CD23, CD23A, CLEC4J, FCE2, IGEBF, FCER2

#### **PRODUCT SPECIFICATION**

# **Molecular Weight**

32 kDa (283aa)

#### **Concentration**

0.25mg/ml (determined by absorbance at 280nm)

#### **Formulation**

Liquid in. Phosphate-Buffered Saline (pH 7.4) containing 10% glycerol

#### **Purity**

> 90% by SDS-PAGE

#### **Endotoxin level**

< 1 EU per 1ug of protein (determined by LAL method)

## **Biological Activity**

Measured by its binding ability in a functional ELISA with Human IgE. The ED50 range  $\leq$  15 ug/ml.

#### Tag

His-Tag

# **Application**

SDS-PAGE, Bioactivity

# **Storage Condition**

Can be stored at +2C to +8C for 1 week. For long term storage, aliquot and store at -20C to -80C. Avoid repeated freezing and thawing cycles.

### **BACKGROUND**



# NKMAXBIO We support you, we believe in your research

# Recombinant human CD23/FCER2 protein

Catalog Number: ATGP4162

# **Description**

CD23/FCER2, also known as low affinity immunoglobulin epsilon Fc receptor isoform a, is a member of subgroup II of the C-type (Ca++-dependent) lectin superfamily. It is a low affinity receptor for B cell specific antigen and IgE. Unlike many of the antibody receptors, CD23 is a C-type lectin. It is found on mature B cells, activated macrophages, eosinophils, follicular dendritic cells, and platelets. It plays an essential role in the growth and differentiation of B cells, and in the regulation of IgE production. This protein also exists in a soluble secretion form and functions as a powerful cleavage-promoting growth factor. Increased levels of soluble CD23/FCER2 lead to the recruitment of unaffected B cells in the presentation of antigen peptides to allergen-specific B cells. Recombinant human CD23/FCER2, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

# **Amino acid Sequence**


<ADP>DTTQSLK QLEERAARNV SQVSKNLESH HGDQMAQKSQ STQISQELEE LRAEQQRLKS QDLELSWNLN GLQADLSSFK SQELNERNEA SDLLERLREE VTKLRMELQV SSGFVCNTCP EKWINFQRKC YYFGKGTKQW VHARYACDDM EGQLVSIHSP EEQDFLTKHA SHTGSWIGLR NLDLKGEFIW VDGSHVDYSN WAPGEPTSRS QGEDCVMMRG SGRWNDAFCD RKLGAWVCDR LATCTPPASE GSAESMGPDS RPDPDGRLPT PSAPLHS<HHH HHH>

#### **General References**

Heyman, B., et al. (2000) Annu. Rev. Immunol. 18:709-737. Bajorath, J., et al. (1996) Protein Sci. 5:240-247.

#### **DATA**

#### **SDS-PAGE**




3ug by SDS-PAGE under reducing condition and visualized by coomassie blue stain.

# **Biological Activity**



# Recombinant human CD23/FCER2 protein

Catalog Number: ATGP4162



Human IgE is coated at 5 ug/ml (100 ul/well) can bind Human CD23/FCER2. The ED50 range  $\leq$  15 ug/ml.

